

# Interactive effects of protein nutrition, genetic growth potential and *Heligmosomoides bakeri* infection pressure on resilience and resistance in mice

JENNIFER C. COLTHERD<sup>1\*</sup>, SIMON A. BABAYAN<sup>2</sup>, LUTZ BÜNGER<sup>3</sup>,  
ILIAS KYRIAZAKIS<sup>1†</sup>, JUDITH E. ALLEN<sup>2</sup> and JOS G. M. HOUDIJK<sup>1</sup>

<sup>1</sup> *Animal Health, SAC, West Mains Road, Edinburgh EH9 3JG, UK*

<sup>2</sup> *Centre for Immunity, Infection & Evolution and Institute of Immunology and Infection Research, Ashworth Laboratories, King's Buildings, West Mains Road, Edinburgh EH9 3JT, UK*

<sup>3</sup> *Sustainable Livestock Systems, SAC, West Mains Road, Edinburgh EH9 3JG, UK*

(Received 7 March 2011; revised 3 May 2011; accepted 10 May 2011)

## SUMMARY

The ability of animals to cope with an increasing parasite load, in terms of resilience and resistance, may be affected by both nutrient supply and demand. Here, we hypothesized that host nutrition and growth potential interact and influence the ability of mice to cope with different parasite doses. Mice selected for high (ROH) or low (ROL) body weight were fed a low (40 g/kg; LP) or high (230 g/kg; HP) protein diet and infected with 0, 50, 100, 150, 200 or 250 L<sub>3</sub> infective *Heligmosomoides bakeri* larvae. ROH-LP mice grew less at doses of 150 L<sub>3</sub> and above, whilst growth of ROH-HP and of ROL mice was not affected by infection pressure. Total worm burdens reached a plateau at doses of 150 L<sub>3</sub>, whilst ROH mice excreted fewer worm eggs than ROL mice. Serum antibodies increased with infection dose and ROH mice were found to have higher parasite-specific IgG1 titres than ROL mice. In contrast, ROL had higher total IgE titres than ROH mice, only on HP diets. The interaction between host nutrition and growth potential appears to differentially affect resilience and resistance in mice. However, the results support the view that parasitism penalises performance in animals selected for higher growth.

Key words: minimum level of parasitism, *Heligmosomoides bakeri*, protein nutrition, growth potential, resilience, resistance, serum antibody, mice.

## INTRODUCTION

Exposure to parasites is ubiquitous in both livestock and wildlife species (Holmes, 1993). In most cases this exposure leads to subclinical infection and immunity is gradually acquired. Development of immunity to gastrointestinal nematodes usually coincides with periods of high growth rates and thus high nutrient requirements (Coop and Kyriazakis, 2001). In addition, parasitic infection can be accompanied by variable pathology (Garside *et al.* 2000) and anorexia (Kyriazakis, 2010), which can readily lead to nutrient scarcity even when animals are fed on good quality foods. Under these conditions, hosts may be forced to allocate scarce nutrient resources between the competing traits of growth and parasite control (Coop and Kyriazakis, 2001). Here, we assess whether this allocation is sensitive to infection pressure and genetic potential for growth.

It has long been proposed that a minimum level of antigenic stimulation or infection pressure may be necessary for stimulation of an immune response upon parasitic infection (Dineen, 1963). In agreement, a certain level of parasite infection pressure is also required to produce clinical infection (Vercruysse and Claerebout, 2001) and this has long been known as the case for *Heligmosomoides bakeri* infections (Ehrenford, 1954). It has been suggested that such 'thresholds' may be lower during periods of nutrient scarcity (Bransby, 1993). In addition, Keymer and Tarlton (1991) observed that the accumulation of *H. bakeri* was proportional to the infection pressure at times of protein scarcity, whereas this pattern of accumulation was not observed at high protein levels.

Rauw *et al.* (1998) suggested that artificial selection for enhanced production traits can lead to detrimental changes in host behaviour, physiology and immunity. These changes may arise from allocation of a higher proportion of scarce environmental resources towards the trait selected for, leaving fewer resources available for investment in other physiological processes that contribute to overall fitness (Rauw *et al.* 1998; Coop and Kyriazakis, 1999; Doeschl-Wilson *et al.* 2008). Consequently, we predict that increased resource availability would

\* Corresponding author: Disease Systems Team, SAC, West Mains Road, Edinburgh EH9 3JG, UK. Tel: +44 (0) 131 5353058. Fax: +44 0 131 5353121. E-mail: jcoltherd@hotmail.com

† Current address: School of Agriculture, Food and Rural Development, Newcastle University, Newcastle upon Tyne NE1 7RU, UK.

minimize these negative consequences of artificial selection for enhanced production traits.

We recently demonstrated that mice selected for high body weight (Roslin-high mice, ROH) were less resilient (measured as performance under infection (described by Albers *et al.* 1987)) and less resistant to a single primary infection of 250 L<sub>3</sub> *H. bakeri* than their low growth counterparts (Roslin low growth mice, ROL), especially when given access to low protein foods (Coltherd *et al.* 2009). This suggests that hosts with a higher potential for productive functions may allocate more nutrients to such functions when nutrients are scarce, at the expense of functions involved with limiting parasitism, including anti-helminth immune responses (Wahid *et al.* 1994). Here, we hypothesize that this penalty on resilience in ROH mice will decrease with lower infection levels and that ROL mice will tolerate higher infection levels. We also hypothesize that the levels of infection, which would be tolerated are lower at times of protein scarcity, and that this nutritional sensitivity would be more pronounced in mice genetically selected for high growth (ROH).

#### MATERIALS AND METHODS

##### *Animals and housing*

A cross of 2 inbred lines (C57BL/6J × DBA/2J) was used as the foundation population for a selection experiment in which divergent selection for high and low body weight (BW) at day 42 of age took place over 20 generations, resulting in due course in ROH and ROL lines, respectively (Heath *et al.* 1995; Bünger *et al.* 2001a). Under maintenance conditions and when offered a standard diet, at 42 and 70 days of age respectively, ROH mice would be expected to reach, on average, weights of 36 g and 41 g, with ROL mice reaching 16 g and 19 g, respectively (Bünger *et al.* 2001b). For this experiment, we obtained 72 ROH and 72 ROL male mice inbred for 40 generations, weaned at 21–23 days and housed individually in a controlled environment with an ambient temperature of 21 ± 1 °C and a 12 h light cycle. At weaning, our ROH mice had a body weight of 15.4 ± 0.25 g, whereas ROL mice weighed 7.4 ± 0.26 g. Mice were housed in solid-bottomed cages with fresh sawdust and bedding material was provided twice weekly. The experiment details described below were approved by the Animal Experiment Committee of Scottish Agricultural College (ED AE 26/2007) and carried out under Home Office regulations (PPL 60/3626).

##### *Diets*

All mice were fed *ad libitum* standard expanded breeding diet (RM3(P), Special Diet Services, Witham, UK; digestible crude oil: 38 g/kg; digestible

Table 1. Ingredients and analysis of the experimental diets

Ingredients (g/kg)	Experimental diets	
	LP	HP
Rice starch	556.8	332.0
Maltodextrin	132.0	132.0
Sucrose	100.0	100.0
Soya oil	70.0	70.0
Cellulose	50.0	50.0
Vitamins, minerals and amino acids	48.2	51.4
Casein edible acid	43.0	264.0
Analysis		
Dry matter (g/kg)	902.00	878.40
Crude protein (g/kg dry matter)	46.70	219.00

crude protein (CP): 202 g/kg; starch: 339 g/kg; sugars: 44 g/kg; digestible energy (DE), 12.2 MJ/kg) for 1 week after arrival. Two iso-energetic experimental diets (15 MJ DE/kg) with a fixed amino acid to CP ratio were used, formulated to provide 40 (LP) and 230 (HP) g CP per kg (Table 1). These CP levels were expected to result in protein scarcity or protein adequacy, respectively, relative to expected requirements in non-infected mice (NRC, 1995) and as informed by our previous work using a wide range of dietary CP levels on these inbred lines (Coltherd *et al.* 2009). As casein was used as the protein source, 15 g of cysteine were added to each kg of casein to account for the scarcity of sulphur-containing amino acids.

##### *Infection protocol and experimental design*

The strain of *H. bakeri* used was obtained from Professor Jerzy Behnke (University of Nottingham, UK), and maintained in our laboratory through passages in C57BL/6 mice (see Jenkins and Behnke, 1977 for full origin details). The parasite corresponds to *H. polygyrus bakeri* reported by Duret-Desset *et al.* (1972). At day 0 of the experiment (see below) mice received a single infection of 0, 50, 100, 150, 200 or 250 *H. bakeri* infective larvae (L<sub>3</sub>) suspended in 0.2 ml of water via oral gavage. The highest level of infection pressure (250 L<sub>3</sub>) reduced growth performance in our previous work (Houdijk and Bünger, 2006, 2007; Coltherd *et al.* 2009).

Mice within each line, infected at one of the 6 experimental infection levels (referred to as 0, 50, 100, 150, 200 and 250), were given *ad libitum* access to either LP or HP foods. The experiment was thus 2 lines × 6 levels of infection × 2 protein diets; each cell within this design contained 6 replicates, thus a total of 144 mice were used. The degree of replication was based on our previous experiment for these mouse lines and this parasite model (Coltherd *et al.* 2009). Figure 1 illustrates the experimental design and




Fig. 1. Diagram of experimental design: timeline in experimental days shown along the bottom and age of mice in parentheses.

timing of procedures. Mice entered the experiment (day 0) through an adaptation phase (day -14 to day 0), which included a period where a 50:50 mix of experimental and standard diet was offered to acclimatize the mice to the experimental diet (day -7 to day -4) followed by feeding the experimental diets alone (day -4 onwards). This timescale of experimental diet feeding treatment allowed the experiment to be carried out whilst mice were still growing. Mice were humanely killed on day 28 post-infection (p.i.), for the assessment of worm burdens, colon egg count, body fat percentage and serum antibodies (see below).

#### Sample measurements and collection

**Body weight and food intake.** Between day 0 and 28 p.i., mice and food refusals were weighed twice weekly (Tuesday and Friday) resulting in 8 experimental periods for food intake. On each of these days food refusals were weighed out and fresh food weighed in. Around 30 g were offered to ROH and 15 g to ROL mice to ensure *ad libitum* feeding (Coltherd *et al.* 2009). Average daily food intake was calculated per day for each of the 8 experimental periods for each mouse, as well as mean food intake over the total 8 experimental periods.

**Nematode egg count in feces.** Mice were placed onto wire-bottomed cages, with access to food and water,

overnight for fecal sample collection on days 21 and 25 p.i. A modified flotation technique (Christie and Jackson, 1982) was used to assess the concentration of nematode eggs/g feces. The total period of fecal collection (finish time – start time) was recorded to calculate an estimated feces volume/12 h/mouse. This value was used to standardize egg output (EO, eggs excreted/12 h) to eliminate the dilution effect expected for ROH mice due to their larger size, food intake and thus larger volumes of feces (Coltherd *et al.* 2009).

**Worm burden and nematode egg count in colon contents.** Mice were killed humanely on day 28 via CO<sub>2</sub> inhalation and dissected to obtain the small intestine and the colon. The small intestine was weighed, opened and placed directly into a 5% formaldehyde solution pending assessment of the number of male and female worms. The colon contents were weighed and a colon egg count (eggs/g) performed for every mouse. The resulting colon egg count was then multiplied by the colon contents weight to eliminate any dilution effects and obtain a final number of eggs in colon (EIC) value. The EIC was divided by the number of females counted to estimate *per capita* fecundity (eggs/female).

**Fat percentage.** To predict percentage of carcass fat, the mouse carcasses were weighed and bagged upon dissection for subsequent freeze-drying. In

preparation for the freeze-drying process, incisions were made, to allow maximal water loss, in the back, tail and head of the animal. When weight loss ceased (approximately 7 days later) the dried carcasses were re-weighed. Fat percentage was calculated as  $((\text{freeze-dried weight} \times 1.13) / \text{carcass weight}) - 0.302 \times 100$  (Hastings and Hill, 1989).

#### Enzyme-linked immunosorbent assays – ELISAs

At dissection, blood was collected from the chest cavity, after severing the vessels around the heart, into a 2 ml vol. tube containing Serasieve® (Hughes and Hughes Ltd, Wellington, UK) and serum collected via centrifugation at 1500 g. Serum IgG1 was measured as a key antibody associated with Th2 responses, whilst IgE was measured as an antibody specifically implicated in protective immunity against parasites like *H. bakeri* (Urban *et al.* 1991; 1992; Wahid *et al.* 1994; Negrao-Correa *et al.* 1999) and specifically regulated by the Th2 cytokine IL-4 (Silva *et al.* 2006). Antibody levels were used as read-outs for an indirect measurement of immune costs to indicate possible trade-offs. The concentration of anti-*H. bakeri* antibody IgG1 was measured by an indirect ELISA. Mouse sera were measured individually. Fifty  $\mu\text{l}$  of *H. bakeri* soluble extract in carbonate buffer (5  $\mu\text{g}/\text{ml}$ ) were added to each well of a 96-well plate and left to incubate overnight at 4 °C. Plates were then washed with PBS-Tween and blocked using 100  $\mu\text{l}$  of PBS + BSA 4% for 1 h at 37 °C in the dark. Plates were again washed with PBS-Tween, 50  $\mu\text{l}$  of serum, serial-diluted 11 times from 1/400 in PBS-BSA 1%, added and the plates incubated overnight at 4 °C. Plates were washed with PBS-Tween and 50  $\mu\text{l}$  of HRP-conjugated detection antibody was used. The plates were finally incubated in 50% TMB – 50% H<sub>2</sub>O<sub>2</sub> (KPL) under silver foil to protect from the light until the top two standards had saturated. Colour development was then stopped using 25  $\mu\text{l}$  of 1 M HCl, and plates were read at 450 nm.

The concentration of total IgE antibody was measured by a capture ELISA. Mouse sera were measured individually. The 96-well plates were coated with 50  $\mu\text{l}$  of IgE (2  $\mu\text{g}/\text{ml}$ ) capture antibody and left at 4 °C overnight. Capture antibody was flicked off and 100  $\mu\text{l}$  of Marvel solution:carbonate buffer (5%) were added and left to incubate for 2 h at 37 °C. Plates were then washed in TBST, a row of standards was created from a top standard of 5  $\mu\text{g}/\text{ml}$  and doubling down across the columns, the sera wells were diluted 1/10 and 1/20 and the plates were incubated at 37 °C for 2 h. Plates were again washed in TBST, 50  $\mu\text{l}$  of biotinylated 2° detection IgE (2  $\mu\text{g}/\text{ml}$ ) added then incubated for a further 1 h at 37 °C. Plates were washed in TBST, 100  $\mu\text{l}$  of conjugate Extravidin Peroxidase (1/8000) added and the plates incubated at 37 °C for 30 min. Plates were

finally washed in TBST, then in H<sub>2</sub>O before 100  $\mu\text{l}$  of substrate TMB were added to each well, and the plate developed in darkness until saturation had occurred, and was then stopped using 100  $\mu\text{l}$  of 1 M HCl. The plate was then read at 450 nm on a spectrophotometer.

#### Statistical analysis

Due to the skewed nature of the data, Egg Output (EO), Eggs In Colon (EIC) and worm burdens were Log<sub>10</sub> (n+1) transformed. To account for the relatively large differences in performance data between the mouse lines, arising from *a priori* differences in body weight, feed intake and body weight gain data (both in mg per day) were also Log<sub>10</sub> transformed before analysis (Falconer and MacKay, 1996; Coltherd *et al.* 2009). Repeated measures through Restricted Maximal Likelihood (REML) was used to assess the interactive effects of line, dietary CP content, infection dose and time on food intake. The interactive effects of line, dietary protein content and infection pressure on mean food intake, body fat percentage, EO, worm burden data and serum antibody titres were analysed with REML to allow inclusion of litter origin as a random effect. Interactions that did not approach significance were omitted from the final model. EO, EIC and total worm burdens were checked for non-linearity by fitting a quadratic regression using infection pressure as an explanatory variable. Significance is indicated by a *P*-value less than 0.05 and where appropriate trends are indicated by a *P*-value between 0.051 and 0.1. All reported error values are standard errors of the mean. All statistical analyses were performed using Genstat 11 for Windows release 11.1, 2008 (Lawes Agricultural Trust, Rothamsted, UK).

## RESULTS

### Food intake and body weight gain

Resilience was measured by assessing food intake and weight gain under infection. Figure 2 shows Log<sub>10</sub>-transformed averaged daily food intake and average daily weight gain. Mouse line and dietary protein content interacted significantly for average daily food intake ( $F_{1,96} = 16.28$ ,  $P < 0.001$ ; Fig. 2a). This interaction arose from ROH-0 mice on the LP food consuming less food than their HP counterparts, whilst in contrast ROL-0, ROL-50 and ROL-100 mice on the LP food consumed more food than their HP counterparts. As expected, ROH mice consumed more food than ROL mice ( $F_{1,32} = 260.20$ ,  $P < 0.001$ ). In addition, there were no 2-, 3- or 4-way interactions containing time and infection pressure ( $P > 0.10$ ) indicating the absence of observable anorexia. However, time did interact with mouse line and dietary protein ( $F_{8,936} = 5.25$ ,




Fig. 2. (a)  $\text{Log}_{10}$  transformed daily food intake and (b) body weight gain of high (ROH – open circles) and low (ROL – closed circles) body weight mice averaged over 28 days fed on a high (solid lines) or low (dashed lines) crude protein diet.

$P < 0.001$ ) for food intake; ROL-LP mice ate more than ROL-HP mice in each period, whilst ROH-LP ate more than ROH-HP during periods 1 and 2 only, but ate less than ROH-HP from period 4 onwards (data not shown).

All mice gained weight during the experiment. ROH mice gained an average of  $0.36 \pm 0.019$  g/d while ROL mice gained on average  $0.15 \pm 0.005$  g/d. Genetic growth and dietary protein interacted for weight gain ( $F_{1,117} = 50.12$ ,  $P < 0.001$ ). This was due to ROH mice reducing their weight gain on the LP diet whilst ROL mice were unaffected. A trend was observed for the 3-way interaction between genetic growth, dietary protein content and infection pressure for mean daily weight gain ( $F_{5,117} = 1.96$ ,  $P = 0.09$ ; Fig. 2b). Weight gain of ROH-LP mice was similar as infection pressure increased from 0 to 100  $L_3$ , but significantly decreased as infection pressure further increased 150 to 200 and 250  $L_3$ , whilst infection pressure did not affect weight gain of ROH-HP and ROL mice. However, in the absence of infection, the effect of dietary protein on weight gain differed between ROH and ROL mice; weight gain of ROL-LP mice was the same as that of ROL-HP mice, whilst weight gain was lower for ROH-LP mice compared to ROH-HP mice ( $F_{1,117} = 86.10$ ,  $P < 0.001$ ).

#### Body fat percentage

The percentage body fat was measured to assess the extent to which excess energy was stored by mice.




Fig. 3. Average body fat percentage, taken 28 days after infection, of high (ROH – open circles) and low (ROL – closed circles) body weight mice fed on a high (solid lines) or low (dashed lines) crude protein diet.

The interaction between mouse line and dietary protein content was highly significant for body fat percentage (Fig. 3;  $F_{1,97} = 14.41$ ,  $P < 0.001$ ); independent of infection pressure, ROL-HP mice had a lower body fat percentage than ROL-LP mice, whilst feeding treatment did not affect fat percentage in ROH mice, causing this interaction to be observed. Parasitism did not impose an energetic requirement that could not be met by the experimental diets fed in this experiment.

#### Egg output (EO), eggs in the colon (EIC) and total worm burden (TWB)

Resistance to parasitism was measured by assessing worm burden and parasite egg output. Figure 4 shows the  $\text{Log}_{10}(n+1)$  transformed mean EO, EIC and TWB. In addition, Table 2 provides back-transformed means with 95% confidence intervals for TWB, male and female burdens and *per capita* fecundity. The mean EO averaged over days 21 and 25 p.i. was analysed as there was no interaction between time and infection dose, dietary protein and/or genetic mouse line ( $P > 0.10$ ). EO, EIC and TWB were found to be non-linear ( $P < 0.05$ ). EO, EIC and TWB increased with increased infection pressure ( $F_{4,83} = 25.93$ ,  $P < 0.001$ ,  $F_{4,98} = 17.72$ ,  $P < 0.001$  and  $F_{4,86} = 42.3$ ,  $P < 0.001$  respectively; Fig. 4) until a plateau was reached at 150  $L_3$ . ROL mice had a higher EO and EIC than ROH mice ( $F_{1,30} = 8.21$ ,  $P = 0.008$  and  $F_{1,98} = 10.73$ ,  $P = 0.001$  respectively). However, mouse line and dietary protein content tended to interact for EO; ROL-HP had higher EO than ROL-LP at the lowest infection pressure only ( $F_{1,83} = 3.26$ ,  $P = 0.075$ ; Fig. 4a). Thus, increasing infection pressure over 150  $L_3$  did not further increase parasite load.

Mouse line and dietary protein content interacted for a number of male worms; ROH-HP mice had




Fig. 4.  $\text{Log}_{10}(n+1)$  transformed: (a) 12 h Egg Output (EO) averaged over days 21 and 25 post-infection; (b) Eggs in the Colon (EIC) on day 28 post-infection and (c) total worm burden on day 28 post-infection for high (ROH – open circles) and low (ROL – closed circles) body weight mice, infected with different levels of *Heligmosomoides bakeri* and fed on a high (solid lines) or low (dashed lines) crude protein diet.

fewer male worms than ROH-LP mice whilst feeding treatment did not affect the number of male worms in ROL mice ( $F_{1,99} = 4.48$ ,  $P = 0.037$ ; Table 2). The 3-way interaction between mouse line, dietary protein content and infection pressure tended towards significance for TWB composition ( $F_{4,94} = 2.26$ ,  $P = 0.069$ ). This was due to a higher percentage of female worms in ROH-HP ( $62.6 \pm 3.97\%$ ) mice across all infection pressures, when compared to ROH-LP ( $45.0 \pm 3.07\%$ ) whilst, in contrast, dietary protein content did not systematically affect TWB composition in ROL mice. *Per capita* fecundity was found to be higher in ROL mice ( $F_{1,93} = 8.31$ ,  $P = 0.005$ ) than in ROH mice but was not affected by either dietary protein contents or infection pressure (Table 2). Sex ratios of *H. bakeri* were altered in ROH mice by an HP diet reducing the numbers of male worms when compared to mice on the LP diet.

### Serum antibody titres

Figure 5 shows the serum IgE and IgG1 titres, which demonstrate that an immune response was induced at all infection levels. Mouse line and dietary protein content interacted for total IgE ( $F_{1,101} = 6.77$ ,  $P = 0.011$ ; Fig. 5a); averaged IgE optical density was higher for ROL than for ROH on HP diets, but were similar for LP diets. Although Fig. 5a suggests that these interactions were observed at the 50 and 100 L3 infection pressure only, interactions between mouse line, dietary protein contents and infection pressure were not significant ( $P > 0.17$ ). However, total IgE increased with increased infection pressure ( $F_{5,102} = 51.82$ ,  $P < 0.001$ ).

Mouse line and infection pressure interacted for anti-worm IgG1 ( $F_{5,90} = 7.82$ ,  $P < 0.001$ ; Fig. 5b). IgG1 titres in ROL mice increased in a dose-dependent manner up to 200L<sub>3</sub>, whilst IgG1 titres were higher in ROH mice ( $F_{1,31} = 14.69$ ,  $P < 0.001$ ), and increased linearly with infection pressure up to 250L<sub>3</sub>.

### DISCUSSION

The results of this study support our hypothesis that selection for increased body weight induces a penalty of parasitism on resilience above a certain level of infection ( $>150\text{L}_3$  *H. Bakeri*) but only when protein supply is scarce. However, contrary to our previous findings (Coltherd *et al.* 2009) and our hypothesis, selection for increased body weight did not detrimentally affect resistance traits, as mice selected for low body weight displayed increased nematode egg excretion. This leads to the suggestion that ROL mice may be more resilient to parasitic infection and can maintain growth despite higher levels of parasitism. On the other hand resilience in ROH mice is sensitive to nutrient scarcity. It remains to be ascertained whether this disparity in resistance between the mouse lines is due to selection for body weight inadvertently co-selecting genes associated with immunity (e.g. different MHC alleles; Behnke and Wahid, 1991). Despite our intention, this study failed to achieve protein scarcity in ROL mice, meaning that some results and comparisons may be confounded by protein-sufficient conditions for all ROL mice being compared with protein-deficient and sufficient ROH mice. For this reason, the main comparisons between mouse lines will be between individuals fed the high protein diet to ensure a similar nutritional status.

### Body weight, food intake and fat percentage

A reduction in body weight gain was observed in ROH mice due to reduced dietary protein content and, during protein scarcity, gain was further reduced when infection pressure increased to a dose

Table 2. Mean total worm burdens, males and females, and *per capita* fecundity (PCF) in dependence of mouse line (ROH *vs* ROL), dietary CP content (LP *vs* HP) and infection pressure (50, 100, 150, 200 or 250 *L*<sub>3</sub> *Heligmosomoides bakeri*) with their 95% confidence intervals (CI)

Line	Diet	Infection	Total	Males	Females	PCF	
ROH <sup>a</sup>	LP <sup>b</sup>	50	10 (5–19)	5 (3–9)	5 (3–8)	127 (98–163)	
		100	12 (6–21)	6 (3–11)	6 (3–10)	151 (120–190)	
		150	76 (50–117)	38 (25–57)	38 (25–59)	169 (136–209)	
		200	78 (55–112)	37 (29–47)	44 (24–79)	113 (78–163)	
		250	97 (75–126)	54 (43–67)	46 (26–81)	105 (87–127)	
	HP <sup>b</sup>	50	9 (6–14)	5 (3–7)	4 (3–7)	143 (93–218)	
		100	6 (4–10)	2 (1–3)	4 (3–7)	67 (42–108)	
		150	29 (23–36)	10 (9–12)	18 (14–24)	128 (117–140)	
		200	58 (45–74)	23 (18–30)	34 (27–44)	108 (86–135)	
		250	95 (64–142)	38 (18–30)	57 (39–82)	91 (75–110)	
	ROL <sup>a</sup>	LP	50	8 (5–13)	4 (2–5)	4 (3–6)	207 (84–508)
			100	20 (13–30)	8 (5–13)	12 (8–17)	250 (196–318)
			150	47 (36–62)	22 (16–30)	25 (19–33)	110 (89–136)
			200	78 (65–95)	38 (30–48)	40 (34–47)	209 (177–247)
			250	119 (90–156)	59 (44–79)	60 (47–77)	144 (129–162)
HP		50	10 (7–13)	3 (2–5)	6 (5–8)	190 (146–247)	
		100	23 (17–30)	11 (9–15)	11 (8–15)	177 (151–208)	
		150	52 (40–67)	25 (18–34)	27 (22–34)	201 (153–263)	
		200	84 (65–109)	40 (30–53)	44 (35–56)	130 (103–163)	
		250	103 (70–151)	46 (32–67)	57 (28–84)	171 (147–199)	
<i>P</i> -values							
Line (L)			0.171	0.186	0.135	0.006	
Diet (D)			0.245	0.026	0.886	0.629	
Infection (I)			<0.001	<0.001	<0.001	0.913	
L × D			0.101	0.011	0.590	0.117	
L × I			0.161	0.082	0.603	0.219	
D × I			0.929	0.964	0.899	0.120	
L × D × I			0.879	0.629	0.762	0.397	

<sup>a</sup> Roslin line mice selected for high (ROH) and low (ROL) 42d body weight.

<sup>b</sup> 30 g crude protein (CP)/kg diet (LP) and 230 g CP/kg diet (HP).

over 150L<sub>3</sub>. In contrast, ROL mice performance was not affected by infection, whilst uninfected ROL-LP mice gained more weight than their HP counterparts. This weight gain in ROL-0-LP was associated with higher feed intake on the LP diet compared to the HP diet, which was likely the result of successful attempts to increase protein intake from a low protein diet (Kyriazakis *et al.* 1991). The resulting excess energy intake would need to be lost as heat or stored as fat

(Emmans and Kyriazakis, 2000); the latter is consistent with the higher body fat percentage observed for ROL-LP mice compared to ROL-HP mice in this and our earlier study (Coltherd *et al.* 2009). There was no effect of infection on body fat percentage in this experiment in contrast to the findings of calorie restrictive and metabolic stress experiments, which show that infection decreases body fat percentage (Kristan and Hammond, 2001, 2006).




Fig. 5. (a) Total serum IgE antibody titres, and (b) anti-*H. bakeri* IgG1 serum titre for high (ROH—open circles) and low (ROL—closed circles) body weight mice fed a high (solid lines) or low (dashed lines) crude protein diet.

Under the theory that nutrients are partitioned towards artificially selected traits (Rauw *et al.* 1998; Coop and Kyriazakis, 2001; Doeschl-Wilson *et al.* 2009) and away from unselected ones, ROH growth on the low protein diet suggests that this pattern of nutrient partitioning may indeed have been the case at low levels of infection. The reduction of body weight gain at higher infection pressures (150L<sub>3</sub> and above) is possibly an outcome of the intensity of the infection being sufficiently high to potentially affect host fitness. Consequently, it would be expected that the host would divert nutrients away from growth functions and towards immune function to limit the pathology and/or fitness consequences caused by parasitism and increase the chances of host survival. Therefore, this suggests a novel hypothesis, i.e. that nutrient allocation is sensitive to infection pressure.

The lack of parasite-induced anorexia across the infected mice was unexpected because such a reduction had been observed in previous studies with this mouse–parasite model (Houdijk and Bünger, 2006; Coltherd *et al.* 2009) and in other studies using *H. bakeri* (Boulay *et al.* 1998; Ing *et al.* 2000; Tu *et al.* 2007). A reduction in feed intake would be expected to occur mainly during the initial stages of infection (Coltherd *et al.* 2009; Kyriazakis, 2010), and it might be that this would only have been evident above a certain level of infection pressure (Sandberg *et al.* 2006). Therefore, it cannot be excluded that diluting effects of mice infected below such a level of infection pressure did not allow us to statistically detect the presence of anorexia. Indeed,

assessing the effect of infection on feed intake at each infection pressure separately showed that only mice infected with 250 L<sub>3</sub> experienced anorexia observed between days 7 and 10 post-infection ( $F_{8,296} = 2.68$ ,  $P = 0.039$ ; data not shown). However, this effect was small, and supported by the absence of a meaningful interaction between infection pressure and time for feed intake when analysing all infection pressures simultaneously ( $F_{40,936} = 1.46$ ,  $P = 0.102$ ).

#### Worm burden and egg output

As infection pressure increased, the worm burden and number of eggs released increased to a maximum level and then appeared to plateau. This maximum level was lower in ROH mice than in ROL mice, which was contrary to our previous findings (Coltherd *et al.* 2009). The observed plateau in egg output and worm burden in ROH mice suggests an optimal parasite load controlled by the host and corresponds to the further reduction in body weight gain seen at 150L<sub>3</sub> and above. Indeed Paterson and Viney (2002) proposed that host immune responses reduced parasite survival to limit parasite load as they saw no effect of infection pressure on establishment of parasites, but did observe that survivorship of parasites during a heavy infection was significantly reduced as duration of infection increased. Density-dependent effects on parasite numbers and egg output are commonly observed. For gastrointestinal parasites this relationship is usually negative (or inverse), i.e. as density of the parasite population increases worm survival and fecundity decrease (Bishop and Stear, 2000; Bleay *et al.* 2007). Christensen *et al.* (1995) found that egg production and *per capita* fecundity progressively reduced as infection pressure of *Oesophagostomum dentatum* increased from 2000 to 20000 and 200000 L<sub>3</sub>. This observation of *per capita* fecundity differs from our results, where infection pressure and dietary protein content did not affect fecundity. Observations made by Keymer and Slater (1987) suggest that *per capita* fecundity is highly variable when worm burden is low, with around 78% of the worms having similar egg production to hosts with higher worm burdens. This observation may be an explanation for the lack of density-dependent effects on fecundity in our model, as we also found high variability in egg counts at lower levels of infection (data not shown). There are 2 main proposed mechanisms for the observation of density-dependent egg production (Anderson and Michel, 1977; Kerboeuf and Jolivet, 1984; Irvine *et al.* 2001). Firstly, intra-specific competition may limit resources for the natural development of adult parasites and the production of eggs by adult females. Secondly, host immune responses may stunt worm development directly, limiting female size and thus fecundity. From the latter explanation it would be

expected that infections in immune animals are more sensitive to nutritional manipulation, as has been shown in several different species (Keymer and Tarlton, 1991; Ing *et al.* 2000; Houdijk *et al.* 2005; Kidane *et al.* 2009). This may account for the lack of evidence of density-dependent effects on egg production during this study using a single-dose primary infection.

The lower proportion of female worms present in ROH-LP TWB when compared to ROH-HP suggests that decreased protein supply may have altered the sex ratio either through altering the gut environment, e.g. causing villi atrophy (Tu *et al.* 2007) which in turn reduces preferred habitat and feeding opportunities (Bansemir and Sukhdeo, 1994, 1996) and may reduce mating opportunities, or alternatively through disproportionate survival of larvae (Stien *et al.* 2005). One might expect that female worms would be most affected by reduced feeding opportunities due to their larger size when compared to male worms (Poulin, 1997). Indeed, reduced worm egg production has been observed in animals fed very low protein diets (Athanasiadou *et al.* 2001; Coltherd *et al.* 2009). It follows that female worm development, and so capacity for egg production, may have been stunted due to limited feeding opportunities. This perhaps was not observed in the current study due to the low protein diet, 40 g CP per kg, not being sufficiently low to produce effects on fecundity which was observed at 30 g CP per kg previously (Coltherd *et al.* 2009).

#### Antibody concentrations

The serum antibody titres, although used as an indirect measure of a resource cost borne by the immune system, can also give some indication of the polarization of the immune system. The Th2 type antibodies IgE and IgG1 were assessed due to their previously reported elevation during parasitic helminth infection (Urban *et al.* 1991; Romagnani, 1991; Bell *et al.* 1992), and specifically in *H. bakeri* infections (Urban *et al.* 1991; Ben-Smith *et al.* 1999; Negrao-Correa *et al.* 1999). IgE levels were higher in ROL than in ROH mice, with both mice lines showing increasing IgE antibody levels with increasing infection pressure, whilst dietary protein contents did not affect IgE production. Whilst this is in agreement with Boulay *et al.* (1998), Ing *et al.* (2000) did find that at day 28 post-primary infection protein malnutrition reduced IgE titres in an *H. bakeri* infection. This may be due to the different inbred mouse strain used (Behnke *et al.* 2006) and/or a longer adaptation period to the experimental diets. Parasite-specific IgG1 levels were found to increase in response to increasing larval pressure for both mouse lines, with the exception that IgG1

concentration levelled off at 200L<sub>3</sub> for ROL mice whilst continuing to increase in ROH mice. ROH mice also had consistently higher IgG1 levels than ROL mice. Again, we did not observe an effect of dietary protein contents for parasite-specific IgG1, which is in accordance with Boulay *et al.* (1998). The increasing parasite-specific antibody response with increasing larval pressure was also observed during an early antibody response to the liver fluke *Opisthorchis viverrini* in hamsters, as the infection became more chronic (6-month infection) however, this relationship inverted and an infection with 25 metacercariae produced higher responses than 50 and 100 metacercariae (Sripa and Kaewkes, 2000). In contrast, a human study using *Necator americanus*, found that early parasite-specific IgG antibody responses were higher in subjects given 10 larvae when compared to 25 and 50 larvae, whilst later IgG (week 13) responses were lower in subjects given 10 larvae when compared to the higher doses. Furthermore early and late IgE responses were higher in subjects given 25 larvae compared to the other doses (Mortimer *et al.* 2006). In this model, selection for increased body weight does not seem to have been detrimental to serum antibody production and, indeed, IgG1 concentrations were actually higher in ROH mice than in ROL mice. The differences in patterns of IgG1 vs IgE in ROH vs ROL mice in this study may be due to the more strict dependence of IgE on IL-4 (Silva *et al.* 2006). Thus the polyclonal IgE response, as measured here, is likely to be a robust measure of 'Th2' bias, while the antigen-specific IgG1 responses may provide the more reliable measure of investment in parasite-specific immunity.

In conclusion, whilst the results of this study do not support our original hypothesis as far as resistance to parasites is concerned, they are consistent with the view that selection on the basis of high body weight may reduce the resilience to pathogen challenge (Rauw *et al.* 1998). This penalty on performance may be sensitive to parasite load and can be overcome to an extent by increased dietary protein contents. Although increased dietary protein content had no effect on resistance to a primary infection consistently throughout our model (Coltherd *et al.* 2009), it has been demonstrated that increased protein nutrition will improve resistance to *H. bakeri* to a greater extent in a challenge infection (Boulay *et al.* 1998; Ing *et al.* 2000). The observation that the ROL mice showed increased resilience but also increased worm fecundity compared to ROH mice was unexpected and requires further investigation. The data potentially imply that resilient animals can be a source of infection to other animals in their environment as they are likely to be overlooked for drenching due to their lack of symptoms, thus increasing the risk to naïve animals from environmental contamination.

## ACKNOWLEDGEMENTS

This work was carried out at the Scottish Agricultural College and University of Edinburgh. Special Diet Services (Witham, UK) supplied the diets. Statistical input during the experimental design and the analysis of the experiment was received from Biomathematics and Statistics Scotland (BIOSS) colleagues, Ian Nevison and Mintu Nath, whose input is gratefully acknowledged. We would also like to thank Bianca Theeruth, John Verth, Ray McInnes, and Moira Stewart for their technical assistance.

## FINANCIAL SUPPORT

Financial assistance was obtained from the Scottish Government Rural and Environmental Research and Analysis Directorate (RERAD).

## REFERENCES

- Albers, G. A., Gray, G. D., Piper, L. R., Barker, J. S., Le Jambre, L. F. and Barger, I. A. (1987). The genetics of resistance and resilience to *Haemonchus contortus* infection in young merino sheep. *International Journal for Parasitology* **17**, 1355–1363.
- Anderson, R. M. and Michel, J. F. (1977). Density-dependent survival in populations of *Ostertagia ostertagi*. *International Journal for Parasitology* **7**, 321–329.
- Athanasiadou, S., Kyriazakis, I., Jackson, F., and Coop, R. L. (2001). The effects of condensed tannins supplementation of foods with different protein content on parasitism, food intake and performance of sheep infected with *Trichostrongylus colubriformis*. *British Journal of Nutrition* **86**, 697–706.
- Bansemir, A. D. and Sukhdeo, M. V. K. (1994). The food resource of adult *Heligmosomoides polygyrus* in the small-intestine. *Journal of Parasitology* **80**, 24–28.
- Bansemir, A. D. and Sukhdeo, M. V. K. (1996). Villus length influences habitat selection by *Heligmosomoides polygyrus*. *Parasitology* **113**, 311–316.
- Behnke, J. M., Mugambi, J. M., Clifford, S., Iraqi, F. A., Baker, R. L., Gibson, J. P. and Wakelin, D. (2006). Genetic variation in resistance to repeated infections with *Heligmosomoides polygyrus bakeri*, in inbred mouse strains selected for the mouse genome project. *Parasite Immunology* **28**, 85–94.
- Behnke, J. M. and Wahid, F. N. (1991). Immunological relationships during primary infection with *Heligmosomoides polygyrus* (*Nematospiroides dubius*): H-2 linked genes determine worm survival. *Parasitology* **103**, 157–164.
- Bell, R. G., Appleton, J. A., Negrao-Correa, D. and Adams, L. S. (1992). Rapid expulsion of *Trichinella spiralis* in adult rats mediated by monoclonal antibodies of distinct IgG isotypes. *Immunology* **75**, 520–527.
- Ben-Smith, A., Wahid, F. N., Lammas, D. A. and Behnke, J. M. (1999). The relationship between circulating and intestinal *Heligmosomoides polygyrus*-specific IgG<sub>1</sub> and IgA and resistance to primary infection. *Parasite Immunology* **21**, 383–395.
- Bishop, S. C. and Stear, M. J. (2000). The use of a gamma-type function to assess the relationship between the number of adult *Teladorsagia circumcincta* and total egg output. *Parasitology* **121**, 435–440.
- Bleay, C., Wilkes, C. P., Paterson, S. and Viney, M. E. (2007). Density-dependent immune responses against the gastrointestinal nematode *Strongyloides ratti*. *International Journal for Parasitology* **37**, 1501–1509.
- Boulay, M., Scott, M. E., Conly, S. L., Stevenson, M. M. and Koski, K. G. (1998). Dietary protein and zinc restrictions independently modify a *Heligmosomoides polygyrus* (Nematoda) infection in mice. *Parasitology* **116**, 449–462.
- Bransby, D. I. (1993). Effects of grazing management practices on parasite load and weight gain of beef cattle. *Veterinary Parasitology* **46**, 215–221.
- Bünger, L., Laidlaw, A., Bulfield, G., Eisen, E. J., Medrano, J. F., Bradford, G. E., Pirchner, F., Renne, U., Schlote, W. and Hill, W. G. (2001a). Inbred lines of mice derived from long-term growth selected lines: unique resources for mapping growth genes. *Mammalian Genome* **12**, 678–686.
- Bünger, L., Renne, U. and Buis, R. C. (2001b). Body weight limits in mice. In *Encyclopedia of Genetics* (ed. Reeve, E. C. R.), pp. 337–360. Fitzroy Dearborn Publishers, London, UK and Chicago, IL, USA.
- Christensen, C. M., Barnes, E. H., Nansen, P., Roepstorff, A. and Slotved, H. C. (1995). Experimental *Oesophagostomum dentatum* infection in the pig: worm populations resulting from single infections with three doses of larvae. *International Journal for Parasitology* **25**, 1491–1498.
- Christie, M. and Jackson, F. (1982). Specific identification of strongyle eggs in small samples of sheep feces. *Research in Veterinary Science* **32**, 113–117.
- Coltherd, J. C., Bünger, L., Kyriazakis, I. and Houdijk, J. G. M. (2009). Genetic growth potential interacts with nutrition on the ability of mice to cope with *Heligmosomoides bakeri* infection. *Parasitology* **136**, 1043–1055.
- Coop, R. L. and Kyriazakis, I. (1999). Nutrition-parasite interaction. *Veterinary Parasitology* **84**, 187–204.
- Coop, R. and Kyriazakis, I. (2001). Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. *Trends in Parasitology* **17**, 325–330.
- Coop, R. L., Sykes, A. R. and Angus, K. W. (1982). The Effect of 3 Levels of Intake of *Ostertagia-Circumcincta* Larvae on Growth-Rate, Food-Intake and Body-Composition of Growing Lambs. *Journal of Agricultural Science* **98**, 247–255.
- Dineen, J. K. (1963). Immunological aspects of parasitism. *Nature, London* **197**, 268–269.
- Doeschl-Wilson, A. B., Brindle, W., Emmans, G. and Kyriazakis, I. (2009). Unravelling the relationship between animal growth and immune response during micro-parasitic infections. *PLoS One* **4**, e7508.
- Doeschl-Wilson, A. B., Vagenas, D., Kyriazakis, I. and Bishop, S. C. (2008). Exploring the assumptions underlying genetic variation in host nematode resistance (Open Access publication). *Genetics Selection Evolution* **40**, 241–264.
- Duret-Desset, M. C., Kinsella, J. M. and Forrester, D. J. (1972). [Arguments in favour of a double origin for nearctic nematodes of the genus *Heligmosomoides* Hall, 1916]. *Annales de Parasitologie humaine et comparée* **47**, 365–382.
- Ehrenford, F. (1954). Effects of dietary protein on the relationship between laboratory mice and the nematode *Nematospiroides dubius*. *Journal of Parasitology* **40**, 486.
- Emmans, G. C. and Kyriazakis, I. (2000). Issues arising from genetic selection for growth and body composition characteristics in poultry and pigs. In *The Challenge of Genetic Change in Animal Production* (ed. Hill, W. G., Bishop, S. C., McQuirk, B., McKay, J. C., Simm, G. and Webb, A. J.), pp. 36–53. British Society of Animal Science, Edinburgh, UK.
- Falconer, D. S. and MacKay, T. F. C. (1996). *Introduction to Quantitative Genetics*, 4th Edn. Longman Scientific and Technical, Harlow, UK.
- Garside, P., Kennedy, M. W., Wakelin, D. and Lawrence, C. E. (2000). Immunopathology of intestinal helminth infection. *Parasite Immunology* **22**, 605–612.
- Hastings, I. M. and Hill, W. G. (1989). A note on the effect of different selection criteria on carcass composition in mice. *Animal Production* **48**, 229–233.
- Heath, S. C., Bulfield, G., Thompson, R. and Keightley, P. D. (1995). Rates of change of genetic-parameters of body-weight in selected mouse lines. *Genetical Research* **66**, 19–25.
- Holmes, P. H. (1993). Interactions between parasites and animal nutrition - the veterinary consequences. *Proceedings of the Nutrition Society* **52**, 113–120.
- Houdijk, J. G. M. and Bünger, L. (2006). Selection for growth increases the penalty of parasitism on growth performance in mice. *Proceedings of the Nutrition Society* **65**, 68A.
- Houdijk, J. G. M. and Bünger, L. (2007). Interactive effects of selection for growth and protein supply on the consequences of gastrointestinal parasitism on growth performance in mice. *Proceedings of the British Society of Animal Science*, 2007 p. 92.
- Houdijk, J. G. M., Jessop, N. S., Knox, D. P. and Kyriazakis, I. (2005). Secondary infection of *Nippostrongylus brasiliensis* in lactating rats is sensitive to dietary protein content. *British Journal of Nutrition* **93**, 493–499.
- Ing, R., Su, Z., Scott, M. E. and Koski, K. G. (2000). Suppressed T helper 2 immunity and prolonged survival of a nematode parasite in protein-malnourished mice. *Proceedings of the National Academy of Sciences, USA* **97**, 7078–7083.
- Irvine, R. J., Stien, A., Dallas, J. F., Halvorsen, O., Langvatn, R. and Albon, S. D. (2001). Contrasting regulation of fecundity in two abomasal nematodes of Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Parasitology* **122**, 673–681.
- Jenkins, S. N. and Behnke, J. M. (1977). Impairment of primary expulsion of *Trichuris-Muris* in mice concurrently infected with *Nematospiroides-Dubius*. *Parasitology* **75**, 71–78.

- Kerboeuf, D. and Jolivet, G.** (1984). *Heligmosomoides polygyrus*: time of anthelmintic treatment and infection parameters in mice exposed to increasing doses of larvae. *Experimental Parasitology* **57**, 307–315.
- Keymer, A. E. and Slater, A. F.** (1987). Helminth fecundity: density dependence or statistical illusion? *Parasitology Today* **3**, 56–58.
- Keymer, A. E. and Tarlton, A. B.** (1991). The population dynamics of acquired immunity to *Heligmosomoides polygyrus* in the laboratory mouse: strain, diet and exposure. *Parasitology* **103**, 121–126.
- Kristan, D. M. and Hammond, K. A.** (2001). Parasite infection and caloric restriction induce physiological and morphological plasticity. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* **281**, R502–R510.
- Kristan, D. M. and Hammond, K. A.** (2006). Effects of three simultaneous demands on glucose transport, resting metabolism and morphology of laboratory mice. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology* **176**, 139–151.
- Kidane, A., Houdijk, J. G. M., Tolkamp, B. J., Athanasiadou, S. and Kyriazakis, I.** (2009). Consequences of infection pressure and protein nutrition on periparturient resistance to *Teladorsagia circumcincta* and performance in ewes. *Veterinary Parasitology* **165**, 78–87.
- Kyriazakis, I.** (2010). Is anorexia during infection in animals affected by food composition? *Animal Feed Science and Technology* **156**, 1–9.
- Kyriazakis, I., Emmans, G. C. and Whittemore, C. T.** (1991). The ability of pigs to control their protein intake when fed in three different ways. *Physiology and Behaviour* **50**, 1197–1203.
- Mortimer, K., Brown, A., Feary, J., Jagger, C., Lewis, S., Antoniak, M., Pritchard, D. and Britton, J.** (2006). Dose-ranging study for trials of therapeutic infection with *Necator americanus* in humans. *American Journal of Tropical Medicine and Hygiene* **75**, 914–920.
- Negrao-Correa, D., Adams, L. S. and Bell, R. G.** (1999). Variability of the intestinal immunoglobulin E response of rats to infection with *Trichinella spiralis*, *Heligmosomoides polygyrus* or *Nippostrongylus brasiliensis*. *Parasite Immunology* **21**, 287–297.
- NRC.** (1995). *Nutrient Requirements of Laboratory Animals*, 4th Edn. National Academy Press, Washington D.C., USA.
- Paterson, S. and Viney, M. E.** (2002). Host immune responses are necessary for density dependence in nematode infections. *Parasitology* **125**, 283–292.
- Poulin, R.** (1997). Covariation of sexual size dimorphism and adult sex ratio in parasitic nematodes. *Biological Journal of the Linnean Society* **62**, 567–580.
- Rauw, W., Kanis, E., Noordhuizen-Stassen, E. N. and Grommers, F. J.** (1998). Undesirable side effects of selection for high production efficiency in farm animals: a review. *Livestock Production Science* **56**, 15–33.
- Romagnani, S.** (1991). Type-1 T-helper and type-2 T-helper cells: functions, regulation and role in protection and disease. *International Journal of Clinical and Laboratory Research* **21**, 152–158.
- Sandberg, F. B., Emmans, G. C. and Kyriazakis, I.** (2006). A model for predicting feed intake of growing animals during exposure to pathogens. *Journal of Animal Science* **84**, 1552–1566.
- Silva, A. S., Cavalcante, L. T., Faquim-Mauro, E. L. and Macedo, M. S.** (2006). Regulation of anaphylactic IgG1 antibody production by IL-4 and IL-10. *International Archives of Allergy and Immunology* **141**, 70–78.
- Sripa, B. and Kaewkes, S.** (2000). Relationship between parasite-specific antibody responses and intensity of *Opisthorchis viverrini* infection in hamsters. *Parasite Immunology* **22**, 139–145.
- Stien, A., Dallimer, M., Irvine, R. J., Halvorsen, O., Langvatn, R., Albon, S. D. and Dallas, J. F.** (2005). Sex ratio variation in gastrointestinal nematodes of Svalbard reindeer; density dependence and implications for estimates of species composition. *Parasitology* **130**, 99–107.
- Tu, T., Koski, K. G., Wykes, L. J. and Scott, M. E.** (2007). Re-feeding rapidly restores protection against *Heligmosomoides bakeri* (Nematoda) in protein-deficient mice. *Parasitology* **134**, 899–909.
- Urban, J. F., Katona, I. M. and Finkelman, F. D.** (1991). *Heligmosomoides-Polygyrus* - Cd4+ But Not Cd8+ T-Cells regulate the IgE response and protective immunity in mice. *Experimental Parasitology* **73**, 500–511.
- Urban, J. F., Madden, K. B., Svetic, A., Cheever, A., Trotta, P. P., Gause, W. C., Katona, I. M. and Finkelman, F. D.** (1992). The importance of Th2-cytokines in protective immunity to nematodes. *Immunological Reviews* **127**, 205–220.
- Vercruysse, J. and Claerebout, E.** (2001). Treatment vs non-treatment of helminth infections in cattle: defining the threshold. *Veterinary Parasitology* **98**, 195–214.
- Wahid, F. N., Behnke, J. M., Grencis, R. K., Else, K. J. and Ben-Smith, A. W.** (1994). Immunological relationships during primary infection with *Heligmosomoides polygyrus* - Th2 cytokines and primary response phenotype. *Parasitology* **108**, 461–471.